

SUPERSYMMETRY BREAKING ON ORBIFOLDS FROM WILSON LINES *

G. V. GERSDORFF AND M. QUIRÓS

*Instituto de Estructura de la Materia (CSIC), Serrano 123,
E-28006-Madrid, Spain.*

Abstract

We consider five dimensional theories compactified on the orbifold $S^1/\mathbb{Z}_2$ and prove that spontaneous local supersymmetry breaking by Wilson lines and by the Scherk-Schwarz mechanism are equivalent. Wilson breaking is triggered by the $SU(2)_R$ symmetry which is gauged in off-shell $N = 2$ supergravity by auxiliary fields. The super-Higgs mechanism disposes of the would-be Goldstinos which are absorbed by the gravitinos to become massive. The breaking survives in the flat limit, where we decouple all gravitational interactions, and the theory becomes softly broken global supersymmetry.

October 2001

*Work supported in part by CICYT, Spain, under contract AEN98-0816, and by EU under contracts HPRN-CT-2000-00152 and HPRN-CT-2000-00148.

1 INTRODUCTION

Supersymmetry and extra dimensions seem to be essential ingredients in any theory that aims to solve the hierarchy problem, makes gravity compatible with quantum mechanics and unifies gravity with the rest of known particle physics interactions: string theory. Moreover, the possibility of lowering the string (M_s) and compactification ($1/R$) scales [1, 2] to values accessible for present and future accelerators (i.e. the TeV range) has triggered an enormous interest in fields theories with large extra dimensions where gravity [3] and/or matter [4] propagate. In particular, in theories with compactification scales in the TeV range where matter propagates the effect of Kaluza-Klein (KK) excitations can be detected by direct [5] or indirect [6] production at present colliders providing an unambiguous signature of extra dimensions [7].

These theories present the exciting feature that some observables are insensitive to the ultraviolet (UV) cutoff M_s [4, 8–11], and thus they are finite and can be considered as predictions of the theory: this happens in particular with the Higgs mass in supersymmetric higher dimensional models, a fact that has opened a strong debate on the subject [13]. In fact summing over all the tower of KK-excitations when they propagate in internal lines is at the origin of the Higgs mass finiteness in the previously mentioned models. It was argued [13] that the presence of a cutoff $\sim M_s$ in the higher dimensional theory should imply cutting off those sums at KK-modes with masses $M^{(n)}$ such that $M^{(n)} \lesssim M_s$, thus generating quadratic divergences that could not be canceled by supersymmetry. However it was subsequently argued that regularizing KK-sums with a sharp cutoff (truncated towers) was not consistent with the symmetries of the higher dimensional theory. In particular if the underlying theory is a string theory (as we are assuming) the couplings of bulk states to localized states are regularized as $\exp\{-(M^{(n)}/M_s)^2/2\}$ [12] and so their interactions smoothly decouple without spoiling the higher dimensional (tower) structure. Using the stringy regularization it was shown in Ref. [9] that the Higgs mass was finite and coincident with the previous calculations. Furthermore the same conclusion was reached using genuine field-theoretical regularization consistent with the symmetries of the higher dimensional theory, in particular with a Pauli-Villars regulator [11]. As expected, physical observables are independent of the regularization procedure provided the regularization does not spoil any symmetry of the theory.

The ultimate reason for the finiteness of the Higgs mass can be traced back to the underlying string theory. In type I/I' strings, matter corresponds to open strings with ends on the brane and thus with KK-modes along longitudinal directions and winding modes along transverse directions. If transverse directions are very large (gravitational-like) the winding modes are extremely heavy and decouple. Moreover in the limit $1/R \ll M_s$ (where R is the radius of the longitudinal directions) string excitations decouple and one recovers the field theory result. This has been shown in explicit examples [14]. If the compact dimensions the brane is wrapped around possess the structure of an orbifold (a possible requirement from chirality) there can be localized states on the orbifold fixed points which do not feel the compact dimensions; they are similar to the twisted states of the heterotic string, required by the anomaly cancellation condition. On the other hand matter in the bulk for heterotic strings possesses both KK and winding modes with respect to compact dimensions and thus the safe procedure is considering field theory

calculation up to the compactification scale and include the string result at that scale as “threshold corrections” [15].

The origin of electroweak and supersymmetry breaking is possibly one of the more poorly known aspects of gauge theories. In theories with compactified extra dimensions one can break gauge symmetry or supersymmetry by imposing different boundary conditions on different fields (Scherk-Schwarz (SS) mechanism [16, 17]). In particular the SS breaking of a gauge symmetry [18] can be seen, if all fields have periodic boundary conditions, as a spontaneous breaking induced by the vacuum expectation value (VEV) of the extra dimensional component of a gauge field, which plays the role of a Higgs boson in four dimensional theories (Wilson line or Hosotani mechanism [19]). In this case, if the electroweak Higgs is associated with a Wilson line the finiteness of the Higgs mass can follow, even in non supersymmetric string or field models [14, 20].

On the other hand, the SS-mechanism has been used to break supersymmetry by imposing non-trivial boundary conditions on fields transforming non-trivially under a global symmetry of the theory [4, 8]. Recently SS-supersymmetry breaking based on the $N = 2$ $SU(2)_R$ global symmetry has been interpreted as coming from the VEV of the auxiliary component field of the radion superfield, and thus being interpreted as spontaneous breaking of supersymmetry, in a five dimensional (5D) theory compactified on an orbifold [21–23]. Finally supersymmetry-breaking induced by brane localized dynamics has been studied in several recent papers [24–26].

In this paper we want to study the nature of the symmetry that protects the softness of the Scherk-Schwarz breaking in higher dimensional theories and the finiteness of the Higgs mass after electroweak symmetry breaking. We consider 5D theories compactified on the orbifold $S^1/\mathbb{Z}_2$, which allows for chiral matter in the massless sector and thus is suitable for phenomenological applications. We prove that local supersymmetry spontaneously broken by Wilson lines is equivalent to SS supersymmetry breaking. In fact the $SU(2)_R$ symmetry is gauged in off-shell $N = 2$ supergravity and 5D $SU(2)_R$ gauge bosons are auxiliary fields in the minimal supergravity multiplet. When the corresponding Wilson lines acquire a VEV, supersymmetry is spontaneously broken and one recovers the SS-mechanism where different boundary conditions are imposed to non-trivial representations of $SU(2)_R$. The would-be Goldstinos are absorbed by the corresponding gravitinos in the unitary gauge (super-Higgs mechanism [28]) and disappear from the physical spectrum of fields. In summary, local $N = 2$ supersymmetry is spontaneously broken by Wilson lines. The breaking survives in the flat limit, where we decouple all gravitational interactions, and the theory becomes softly broken global supersymmetry. No new counterterms are created by the spontaneous breaking of the gauge (and super) symmetry, which explains the finiteness of the Higgs mass under radiative corrections provided the theory does not contain anomalous $U(1)$ gauge groups ¹.

¹In the presence of a $U(1)$ gauge symmetry one has to take into account a possible Fayet-Iliopoulos (FI) term on the brane which would contribute to the value of the Higgs mass. Since this is not an F-term it is not protected by the supersymmetric non-renormalization theorems but only by the vanishing of the trace of the $U(1)$ charge matrix Y . In theories with $\text{tr } Y \neq 0$ the FI-term is quadratically sensitive to the UV cutoff [27] and thus must be fine-tuned in order to give a small Higgs mass. In theories with $\text{tr } Y = 0$ it is not renormalized and can be consistently set to zero.

2 WILSON LINES VS. SCHERK-SCHWARZ BREAKING

In theories with extra dimensions compactified on a torus or an orbifold, a symmetry can be broken by two mechanisms which are not present in simply connected spaces: the Scherk-Schwarz and the Hosotani (Wilson line) mechanisms. In the Scherk-Schwarz mechanism [16, 17], fields possess non-trivial boundary conditions and are multiple-valued along the extra dimension. Let us assume that a global or local symmetry with generator $\mathcal{Q}$ is used to twist some of the fields under $2\pi R$ -translations:

$$\phi(x^\mu, x^5 + 2\pi R) = e^{2\pi i \omega \mathcal{Q}} \phi(x^\mu, x^5) . \quad (1)$$

This periodicity condition is satisfied if the fields are given by ²

$$\phi(x^\mu, x^5) = e^{i\omega \mathcal{Q} x^5 / R} \tilde{\phi}(x^\mu, x^5) , \quad (2)$$

where $\tilde{\phi}(x^\mu, x^5 + 2\pi R) = \tilde{\phi}(x^\mu, x^5)$. The symmetry generated by $\mathcal{Q}$ is then broken at tree level by the kinetic term.

If the symmetry with $\mathcal{Q}$ -generator is a local one it is possible to break it by giving a VEV to an extra component of the corresponding gauge field A_M ($M = \mu, 5$) [19] (Hosotani or Wilson line mechanism). All fields are periodic in this picture and gauge transformations which preserve this periodicity (the ones with periodic parameters) divide the Wilson lines into equivalence classes which represent the possible vacua of the theory. We can easily label these vacua by the constant configuration of each equivalence class. We will assume from now on a constant $\langle A_5 \rangle$. All non-singlet fields will receive a mass-shift relative to their KK-value through their covariant derivative. For instance, for a flat compact direction the mass of KK-modes $\phi^{(n)}$ is given by

$$\mathcal{M}_{\phi^{(n)}} = \frac{n}{R} \mathbf{1}_G + \langle A_5 \rangle , \quad (3)$$

with $\mathbf{1}_G$ the identity in the fundamental representation of the gauge group G (for non-constant VEVs the mass matrix will not be block-diagonal). It is possible to switch to the Scherk-Schwarz picture by allowing for gauge transformations with non-periodic parameters. By choosing a transformation as

$$\exp \{ i \langle A_5 \rangle x^5 \} \quad (4)$$

to transform away the VEV we end up with non-periodic fields as in Eq. (1), with $\omega \mathcal{Q} \equiv R \langle A_5 \rangle$.

3 FIVE-DIMENSIONAL $N = 2$ OFFSHELL SUPERGRAVITY

To achieve supersymmetry-breaking one usually applies the SS-mechanism to the automorphism group $SU(2)_R$ of the $N = 2$ supersymmetry in five dimensions. It would be desirable to have the Hosotani picture in this case as well. The softness of the SS-breaking

²It has been pointed out [25, 29] in the context of brane-induced supersymmetry breaking that there exist a more general class of solutions to Eq. (1).

(i.e. no additional counterterms except the ones allowed by unbroken supersymmetry) could then easily be explained by the fact that Hosotani breaking is spontaneous. Steps in this direction have been recently taken [22, 23] by giving a VEV to the F -term of the radion field: when coupling the radion superfield to matter charged under $SU(2)_R$ the SS spectrum is reproduced. However the final proof that such a breaking is related to the Hosotani mechanism, and therefore equivalent to SS, requires the introduction of the gravitational sector.

To this end we will consider the recently found off-shell version of $5D$ $N = 2$ supergravity [30, 31]. The complete off-shell particle content together with the orbifold-parity assignments are displayed in table 1. Choices for parity assignments are discussed in Ref. [32]. They have to be consistent with the symmetries of the bulk action and are thus mainly dictated by $N = 2$ supersymmetry and 5D Poincaré invariance. All fields but g_{MN} , ψ_M and A_M are auxiliary only. The fermions are of the symplectic Majorana type and carry an $SU(2)_R$ index. They can be decomposed into two component spinors as

$$\lambda^i = \begin{pmatrix} \lambda_L^i \\ \epsilon^{ij} \bar{\lambda}_{Lj} \end{pmatrix} \quad (5)$$

where we use the convention $\epsilon^{12} = +1$. Frequently we will suppress the $SU(2)_R$ index. The standard procedure to construct the theory is covariantizing the globally supersymmetric transformation rules with respect to local Lorentz and local $SU(2)_R$ transformations and then supercovariantizing with respect to local supersymmetric transformations. After the total covariantization the supersymmetric transformations include covariant derivatives of component fields and of the supersymmetric parameter $\xi^i(x^M)$, as well as additional terms of $\mathcal{O}(\kappa)$ and higher to close the algebra³. The complete expression for these transformations have been given in Ref. [30]. It will be particularly relevant for our purposes the transformation of the gravitino ψ_M which contains the covariant derivative of the transformation parameter:

$$\delta_\xi \psi_M = \frac{1}{\kappa} \mathcal{D}_M \xi + \dots \quad (6)$$

Here $\mathcal{D}_M$ is the covariant derivative with respect to local Lorentz and local $SU(2)_R$ transformations

$$\mathcal{D}_M = D_M + i\kappa \vec{V}_M \vec{\tau} \ , \quad (7)$$

while D_M is the covariant derivative with respect to local Lorentz transformations. The auxiliary field V_M has mass dimension $5/2$ as shown in table 1. Notice that the local supersymmetric transformations, e.g. in Eq. (6), are consistent with the orbifold action if we define $\xi_L^1(x^M)$ as an even function and $\xi_L^2(x^M)$ as an odd one. Together with the parities of the covariant derivatives, i.e. $\mathcal{D}_\mu$ even and $\mathcal{D}_5$ odd, this gives equal parities on both sides of Eq. (6).

Moreover it has been emphasized that the minimal multiplet is too small to construct a physical action out of it [30]. It has to be extended by a matter multiplet with $8_B + 8_F$ components to complete the $48_B + 48_F$ components that a supergravity multiplet must

³We denote by κ the five dimensional gravitational coupling which has mass dimension $-3/2$.

field	description	dim	states	even	odd
g_{MN}	graviton	0	$(15 - 5)_B$	$g_{\mu\nu}, g_{55}$	$g_{\mu 5}$
ψ_M	gravitino	2	$(40 - 8)_F$	$\psi_{\mu L}^1, \psi_{5L}^2$	$\psi_{\mu L}^2, \psi_{5L}^1$
B_M	graviphoton	3/2	$(5 - 1)_B$	B_5	B_μ
$\vec{V}_M$	$SU(2)_R$ gauge field	5/2	$(15 - 3)_B$	$V_\mu^3, V_5^{1,2}$	$V_\mu^{1,2}, V_5^3$
v^{AB}	antisymmetric tensor	5/2	10_B	$v^{\alpha 5}$	$v^{\alpha\beta}$
$\vec{t}$	$SU(2)_R$ triplet	5/2	3_B	$t^{1,2}$	t^3
C	real scalar	7/2	1_B	C	
ζ	$SU(2)_R$ spinor	3	8_F	ζ_L^1	ζ_L^2

Table 1: The field content of the minimal supergravity multiplet in five dimensions which has $40_B + 40_F$ components.

have. One possibility is adding the non-linear multiplet with field content: ϕ_α^i , a scalar where the index α transforms as a doublet of a global $SU(2)$; φ a singlet scalar; V_M , a singlet vector, and; χ^i , an $SU(2)_R$ doublet fermion. The supersymmetric transformations for the non-linear multiplet components were given in Ref. [30] where it is shown that closure of the algebra on φ and V_M requires a supersymmetric constraint between the components of the minimal and non-linear multiplets which removes the extra bosonic component, with respect to the fermionic ones, which appears in the non-linear multiplet. After doing that the kinetic Lagrangian for $SU(2)_R$ doublets, i.e. the 5D gravitino ψ_M^i in the minimal supergravity multiplet, gauginos λ^i in 5D vector multiplets and scalars A^i in hypermultiplets can be written as

$$\mathcal{L}_{\text{kin}} = -\frac{i}{2}\bar{\psi}_M\gamma^{MNP}\mathcal{D}_N\psi_P + \frac{i}{2}\text{Tr}(\bar{\lambda}\gamma^M\mathcal{D}_M\lambda) - \text{Tr}|\mathcal{D}_MA|^2, \quad (8)$$

where the covariant derivatives are given in (7). Full actions for pure supergravity and supergravity coupled to $N = 2$ matter multiplets can be found in Refs. [30, 31].

4 WILSON BREAKING OF LOCAL SUPERSYMMENTRY

To perform a Wilson line breaking we have to consider supergravity since local $SU(2)_R$ requires local supersymmetry. However we will see that at the end one can consistently decouple all gravity interactions by taking the limit $\kappa \rightarrow 0$ while retaining the effect of supersymmetry breaking. In the minimal realization of 5D $N = 2$ supergravity described above the $SU(2)_R$ symmetry becomes local but it is gauged only off-shell, i.e. the corresponding gauge fields $\vec{V}_M$ are auxiliary fields. Upon orbifolding some of the gauge bosons are even, V_μ^3 , and the others, $V_\mu^{1,2}$, odd and so happens with the corresponding gauge transformations,

$$U = \exp(i\vec{\Lambda}\vec{\tau}), \quad (9)$$

where Λ^3 is even and $\Lambda^{1,2}$ odd. This guarantees that the $\mathbb{Z}_2$ parity defining the orbifold in table 1 is stable under gauge transformations. In this environment one can use the

triplet of scalars $\vec{V}_5$ to break $SU(2)_R$ by Wilson lines. As presented in table 1, the fifth components of the gauge bosons $\vec{V}_5$ should have opposite parities: V_5^3 is odd and $V_5^{1,2}$ are even. These assignments are also consistent with the requirement that the supercovariant derivatives (7), which include local $SU(2)_R$ transformations have a well defined parity. Notice that the parity operator acting on $SU(2)_R$ doublets is proportional to τ^3 , which anticommutes with $\tau^{1,2}$. As a consequence $\tau^{1,2}$ in (7) make, when acting on doublets, a parity flip required for consistency of supercovariant derivatives. In particular, there is contribution from the even scalars $V_5^{1,2}$ to the odd derivative $\mathcal{D}_5$ in (7).

Now we would like to turn a (constant) VEV on $\vec{V}_5$ in order to get nontrivial Wilson lines. Observe that by five dimensional gauge invariance there is no potential for $\vec{V}_5$ which is thus a flat direction. Our choice for the parity of the fields allows constant VEVs only for V_5^1 and V_5^2 . Without loss of generality we can choose V_5^2 and require

$$\langle V_5^2 \rangle = \frac{\omega}{\kappa R}. \quad (10)$$

This parametrization gives the correct dimension and also the right behaviour in the decompactification and flat limits. The factor $1/R$ in (10) guarantees that the VEV vanishes in the limit $R \rightarrow \infty$, where the compact dimension becomes infinite, since the corresponding Wilson breaking should disappear in the limit of “non-compact” (infinite) extra dimension. On the other hand, sending $\kappa \rightarrow 0$ we will see that the gravitino mass remains constant as desired. The dimensionless factor ω is identified with the SS-parameter.

The Goldstino is found from the local supersymmetric transformations and it is provided by the fifth component of the gravitino, ψ_5 , which has a nonzero transformation in the vacuum:

$$\delta_\xi \psi_5 = \frac{1}{\kappa} \mathcal{D}_5 \xi. \quad (11)$$

Let us analyze the corresponding super-Higgs [28] effect. The kinetic terms for the gravitino can be decomposed in four-dimensional and extra components as:

$$\begin{aligned} -\frac{i}{2} \bar{\psi}_M \gamma^{MNR} \mathcal{D}_N \psi_R &= -\frac{i}{2} \bar{\psi}_\mu \gamma^{\mu\nu\rho} \mathcal{D}_\nu \psi_\rho + \frac{i}{2} \bar{\psi}_\mu \gamma^{\mu\nu} \gamma^5 \mathcal{D}_5 \psi_\nu \\ &\quad -\frac{i}{2} \bar{\psi}_\mu \gamma^{\mu\nu} \gamma^5 \mathcal{D}_\nu \psi_5 - \frac{i}{2} \bar{\psi}_5 \gamma^{\mu\nu} \gamma^5 \mathcal{D}_\mu \psi_\nu. \end{aligned} \quad (12)$$

We now do the redefinition ⁴

$$\psi_\mu = \psi'_\mu + \mathcal{D}_\mu (\mathcal{D}_5)^{-1} \psi_5, \quad (13)$$

which can be seen as a local supersymmetry-transformation with parameter $\kappa(\mathcal{D}_5)^{-1} \psi_5$ gauging ψ_5 away. We arrive at a “unitary gauge” where ψ_5 has been “eaten” by the 4D gravitino ψ_μ :

$$-\frac{i}{2} \bar{\psi}_M \gamma^{MNR} \mathcal{D}_N \psi_R = -\frac{i}{2} \epsilon^{\mu\nu\rho\sigma} \bar{\psi}'_\mu \gamma_\sigma \gamma^5 \mathcal{D}_\nu \psi'_\rho + \frac{i}{2} \bar{\psi}'_\mu \gamma^{\mu\nu} \gamma^5 \left(\partial_5 + i \frac{\omega}{R} \tau^2 + \dots \right) \psi'_\nu. \quad (14)$$

⁴For constant non-zero ω the operator $\mathcal{D}_5$ is non-singular.

The second term provides the mass term for ψ'_μ , which can be expressed in two component spinors (now omitting any '):

$$\mathcal{L}_{\text{mass}} = \frac{1}{2} (\psi_{\mu L}^1 \ \psi_{\mu L}^2) \sigma^{\mu\nu} \begin{pmatrix} \omega/R & i p_5 \\ -i p_5 & \omega/R \end{pmatrix} \begin{pmatrix} \psi_{\nu L}^1 \\ \psi_{\nu L}^2 \end{pmatrix} + h.c. \quad (15)$$

Here $p_5 \equiv i \partial_5$ is the component of the momentum along the extra dimension and – for a flat extra dimension – takes the values n/R , where n is a non-negative integer. We can see that the flat direction $\langle V_5^2 \rangle$ breaks supersymmetry as in the no-scale models [33] where the gravitino zero mode mass is $m_{3/2} = \omega/R$.

Note that Eq. (15) holds even for x_5 -dependent ω provided that the transformation (13) is nonsingular. In particular, making the extreme choice

$$\omega \propto \delta(x_5) P_0 + \delta(x_5 - \pi R) P_\pi, \quad (16)$$

where P_0 and P_π are the constants parametrizing superpotential VEVs in Ref. [25], we can also incorporate in our off-shell formulation brane-induced supersymmetry breaking and Eqs. (9) and (11) of [25] are reproduced. The main task would then be to diagonalize the mass matrix as done there. The case $P_0 = -P_\pi$ results in trivial Wilson lines and supersymmetry remains unbroken. A possible physical meaning for the choice (16) was given in Refs. [24–26] where it was interpreted as the result of some integrated out brane fields leaving a constant superpotential on each brane ⁵.

The described formalism is of course equally well applicable if one compactifies on S^1 instead of $S^1/\mathbb{Z}_2$. Five-dimensional $N = 2$ local supersymmetry is directly broken down to $N = 0$ by giving a VEV to V_5^i .

Writing the kinetic terms for the gauginos and the hyperscalars explicitly we get

$$\begin{aligned} \frac{i}{2} \bar{\lambda} \gamma^M \mathcal{D}_M \lambda &= \frac{i}{2} \bar{\lambda} \gamma^\mu \mathcal{D}_\mu \lambda + \frac{i}{2} \bar{\lambda} \gamma^5 \left(\partial_5 + i \frac{\omega}{R} \tau^2 + \dots \right) \lambda, \\ A^\dagger \mathcal{D}^2 A &= A^\dagger \mathcal{D}^\mu \mathcal{D}_\mu A - A^\dagger \left(\partial_5 + i \frac{\omega}{R} \tau^2 + \dots \right)^2 A, \end{aligned} \quad (17)$$

while the corresponding superpartners are inert under $SU(2)_R$. The second terms provide the shifted masses for gauginos and hyperscalars. For gauginos the mass Lagrangian can be cast as

$$\mathcal{L}_{\text{mass}} = \frac{1}{2} (\lambda_L^1 \ \lambda_L^2) \begin{pmatrix} \omega/R & i p_5 \\ -i p_5 & \omega/R \end{pmatrix} \begin{pmatrix} \lambda_L^1 \\ \lambda_L^2 \end{pmatrix} + h.c. \quad (18)$$

Similarly for the hyperscalars one obtains,

$$\mathcal{L}_{\text{mass}} = - \begin{pmatrix} A_1^\dagger & A_2^\dagger \end{pmatrix} \begin{pmatrix} p_5^2 + (\omega/R)^2 & i 2\omega p_5/R \\ -i 2\omega p_5/R & p_5^2 + (\omega/R)^2 \end{pmatrix} \begin{pmatrix} A^1 \\ A^2 \end{pmatrix}. \quad (19)$$

Mass matrices for (15), (18) and (19) give rise to the mass eigenvalues $\omega/R \pm p_5$, identical to those obtained in SS-broken theories [4, 8].

⁵Terms proportional to $\delta(x_5 - x_5^{\text{brane}})$ can generally be created by loop effects of bulk and brane couplings [34]. Consideration of loop corrections is beyond the scope of the present paper.

It is easy to see looking at Eq. (17) that we still have supersymmetry breaking in the global limit $\kappa \rightarrow 0$, $m_{3/2}$ fixed. Taking this limit we decouple all gravity degrees of freedom and are left with $N = 1$ global supersymmetry which is broken down to $N = 0$ whenever we have nonzero ω . In the case of a circle, going to the flat limit results in $N = 2$ global supersymmetry broken down to $N = 0$. One can also see some connection of the Hosotani/SS-mechanism with the radion mediation of Refs. [22, 23]. By looking at the linearized (global) transformations one finds that the radion multiplet ⁶


$$(h_{55} + iB_5, \psi_{5L}^2, V_5^1 + iV_5^2 + 4i(t^1 + it^2))$$

transforms separately under global supersymmetry. The auxiliary field of the radion contains explicitly $V_5^1 + iV_5^2$, so the F -term of this superfield will get a VEV as well. In the local theory it induces spontaneous breaking of supersymmetry, but in the global limit the would-be Goldstino is decoupled and the breaking of supersymmetry is soft. Finally the radion VEV remains undetermined at this level by the no-scale nature of the SS-breaking.

5 CONCLUSIONS

We conclude that supersymmetry breaking is mediated already at tree level from the gravity to the matter/gauge sector through the local $SU(2)_R$ couplings. In complete analogy to the case of an ordinary gauge symmetry, the equivalence to SS breaking can be seen by transforming away the VEV of V_5^2 using a multivalued gauge transformation. Finally the softness of the SS-mechanism is explained by the fact that Hosotani breaking is spontaneous. Only counterterms which are allowed by the underlying supersymmetry can appear in the Lagrangian and so the usual non-renormalization theorems apply. This confirms recent results found in explicit one and two loop calculations [9–11].

We summarize the supersymmetry breaking scheme in the following diagram. We denote by $(N = 0)_{\text{local, global}}$ the breakdown of the corresponding $(N \neq 0)_{\text{local, global}}$.



Starting from a genuine 5D $N = 2$ locally supersymmetric theory we compactify on the orbifold $S^1/\mathbb{Z}_2$ thus creating two branes with local $N = 1$ supersymmetry. In the bulk we retain $N = 2$ local supersymmetry with a $\mathbb{Z}_2$ constraint on the parameter ξ , i.e. ξ_L^1 is even and ξ_L^2 odd. Giving a VEV to V_5^2 breaks both supersymmetries spontaneously, the corresponding Goldstino ψ_5 providing the longitudinal components of the 4D gravitino ψ_μ . This mechanism is equivalent to SS-breaking by means of a nonperiodic $SU(2)_R$

⁶ h_{MN} denotes the linearized metric $g_{MN} = \eta_{MN} + \kappa h_{MN}$.

transformation. Taking $\kappa \rightarrow 0$ one arrives then at the SS-broken theories widely considered in the literature [4, 8]. If SS/Wilson breaking is absent, one obtains in the flat limit $N = 1$ supersymmetry for the KK-modes (left vertical arrow). The counterintuitive reduction to $N = 1$ is explained by the fact that the odd parameter ξ_L^2 vanishes in the global limit, while the linearity of the global transformation laws ensures that supersymmetry is realized mode by mode [35, 36].

ACKNOWLEDGMENTS

We thank A. Delgado for useful discussions and for participating at the early stages of this work. The work of GG was supported by the DAAD.

References

- [1] J. Lykken, *Phys. Rev.* **D54** (1996) 3693; G. Shiu and S.-H.H. Tye, *Phys. Rev.* **D58** (1998) 106007; Z. Kakushadze and S.-H.H. Tye, *Phys. Rev.* **D58** (1998) 126001.
- [2] I. Antoniadis, *Phys. Lett.* **B246** (1990) 377; I. Antoniadis, C. Muñoz and M. Quirós, *Nucl. Phys.* **B397** (1993) 515; I. Antoniadis and K. Benakli, *Phys. Lett.* **B326** (1994) 69; I. Antoniadis, K. Benakli and M. Quirós, *Phys. Lett.* **B331** (1994) 313; I. Antoniadis and M. Quirós, *Phys. Lett.* **B392** (1997) 61.
- [3] G.F. Giudice, R. Rattazzi and J.D. Wells, *Nucl. Phys.* **B544** (1999) 3; E.A. Mirabelli, M. Perelstein and M.E. Peskin, *Phys. Rev. Lett.* **82** (1999) 2236; T. Han, J. Lykken and R.-J. Zhang, *Phys. Rev.* **D59** (1999) 105006; J.L. Hewett, *Phys. Rev. Lett.* **82** (1999) 4765.
- [4] A. Pomarol and M. Quirós, *Phys. Lett.* **B438** (1998) 255; I. Antoniadis, S. Dimopoulos, A. Pomarol and M. Quirós, *Nucl. Phys.* **B544** (1999) 503; A. Delgado, A. Pomarol and M. Quirós, *Phys. Rev.* **D60** (1999) 095008.
- [5] I. Antoniadis, K. Benakli and M. Quirós, *Phys. Lett.* **B460** (1999) 176; E. Accomando, I. Antoniadis and K. Benakli, *Nucl. Phys.* **B579** (2000) 3.
- [6] M. Masip and A. Pomarol, *Phys. Rev.* **D60** (1999) 096005; T.G. Rizzo and J.D. Wells, *Phys. Rev.* **D61** (2000) 016007; A. Delgado, A. Pomarol and M. Quirós, *JHEP* **0001** (2000) 030.
- [7] I. Antoniadis and K. Benakli, *Int. J. Mod. Phys.* **27** (2000) 4237.
- [8] R. Barbieri, L.J. Hall and Y. Nomura, *Phys. Rev.* **D63** (2001) 105007; N. Arkani-Hamed, L.J. Hall, Y. Nomura, D. Smith and N. Weiner, *Nucl. Phys.* **B605** (2001) 81; A. Delgado, and M. Quirós, *Nucl. Phys.* **B607** (2001) 99.
- [9] A. Delgado, G.v. Gersdorff, P. John and M. Quirós, *Phys. Lett.* **B517** (2001) 445.
- [10] R. Contino and L. Pilo, [hep-ph/0104130](#); H.D. Kim, [hep-th/0109101](#).

- [11] R. Contino, L. Pilo, R. Rattazzi and A. Strumia, *JHEP* **0106** (2001) 005; Y. Nomura, [hep-ph/0105113](#); N. Weiner, [hep-ph/0106021](#); R. Barbieri, L.J. Hall and Y. Nomura, [hep-th/0107004](#); A. Delgado, G.v. Gersdorff and M. Quirós, *Nucl. Phys.* **B613** (2001) 49; V.D. Clemente, S.F. King and D.A. Rayner, [hep-ph/0107290](#); V.D. Clemente and Y.A. Kubyshin, [hep-th/0108117](#).
- [12] S. Hamidi and C. Vafa, *Nucl. Phys.* **B279** (1987) 465; L. Dixon, D. Friedan, E. Martinec and S. Shenker, *Nucl. Phys.* **B282** (1987) 13; I. Antoniadis and K. Benakli, *Phys. Lett.* **B326** (1994) 69; I. Antoniadis, K. Benakli and A. Laugier, [hep-th/0011281](#).
- [13] D. Ghilencea and H.P. Nilles, *Phys. Lett.* **B507** (2001) 327; H.D. Kim, [hep-ph/0106072](#).
- [14] I. Antoniadis, K. Benakli and M. Quirós, *Nucl. Phys.* **B583** (2000) 35; I. Antoniadis, K. Benakli and M. Quirós, [hep-th/0108005](#).
- [15] D. Ghilencea, H.P. Nilles and S. Stieberger, [hep-th/0108183](#).
- [16] J. Scherk and J.H. Schwarz, *Phys. Lett.* **B82** (1979) 60; *Nucl. Phys.* **B153** (1979) 61; P. Fayet, *Phys. Lett.* **B159** (1985) 121; *Nucl. Phys.* **B263** (1986) 649.
- [17] R. Rohm, *Nucl. Phys.* **B237** (1984) 553; C. Kounnas and M. Porrati, *Nucl. Phys.* **B310** (1988) 355; S. Ferrara, C. Kounnas, M. Porrati and F. Zwirner, *Nucl. Phys.* **B318** (1989) 75; C. Kounnas and B. Rostand, *Nucl. Phys.* **B341** (1990) 641; K. Benakli, *Phys. Lett.* **B386** (1996) 106; C.A. Scrucca and M. Serone, [hep-th/0107159](#).
- [18] A. Masiero, C.A. Scrucca, M. Serone and L. Silvestrini, [hep-ph/0107201](#); L.J. Hall, H. Murayama and Y. Nomura, [hep-th/0107245](#).
- [19] Y. Hosotani, *Phys. Lett.* **B126** (1983) 309; *Phys. Lett.* **B129** (1983) 193; *Ann. Phys.* **190** (1989) 233.
- [20] S. Randjbar-Daemi, A. Salam and J. Strathdee, *Phys. Lett.* **B124** (1983) 345 [Erratum: *Phys. Lett.* **B144** (1983) 455]; H. Hatanaka, T. Inami and C. S. Lim, *Mod. Phys. Lett.* **A13** (2601) 1998; H. Hatanaka, *Prog. Theor. Phys.* **102** (407) 1999; G. R. Dvali, S. Randjbar-Daemi and R. Tabbash, [hep-ph/0102307](#).
- [21] Z. Chacko and M.A. Luty, *JHEP* **0105** (2001) 067; T. Kobayashi and K. Yoshioka, *Phys. Rev. Lett.* **85** (2000) 5527.
- [22] D. Martí and A. Pomarol, [hep-th/0106256](#).
- [23] D.E. Kaplan and N. Weiner, [hep-ph/0108001](#).
- [24] K. A. Meissner, H. P. Nilles and M. Olechowski, *Nucl. Phys.* **B561** (1999) 30.
- [25] J. Bagger, F. Feruglio and F. Zwirner, [hep-th/0108010](#).
- [26] T. Gherghetta and A. Riotto, [hep-th/0110022](#).

- [27] D. M. Ghilencea, S. Groot Nibbelink and H. P. Nilles, [hep-th/0108184](#); R. Barbieri, L. J. Hall and Y. Nomura, [hep-ph/0110102](#).
- [28] E. Cremmer, B. Julia, J. Scherk, P. van Nieuwenhuizen, S. Ferrara and L. Girardello, *Phys. Lett.* **B79** (1978) 231.
- [29] J. A. Bagger, F. Feruglio and F. Zwirner, [hep-th/0107128](#).
- [30] M. Zucker, *Nucl. Phys.* **B570** (2000) 267; *JHEP* **08** (2000) 016; *Phys. Rev.* **D64** (2001) 024024.
- [31] T. Kugo and K. Ohashi, *Prog. Theor. Phys.* **104** (2000) 835, *Prog. Theor. Phys.* **105** (2001) 323; T. Fujita, T. Kugo and K. Ohashi, [hep-th/0106051](#).
- [32] E. Bergshoeff, R. Kallosh and A. Van Proeyen, *JHEP* **0010** (2000) 033.
- [33] E. Cremmer, S. Ferrara, C. Kounnas and D. V. Nanopoulos, *Phys. Lett.* **B133** (1983) 61; A. B. Lahanas and D. V. Nanopoulos, *Phys. Rep.* **145** (1987) 1.
- [34] H. Georgi, A. K. Grant and G. Hailu, *Phys. Rev.* **D63** (2001) 064027; H. Georgi, A. K. Grant and G. Hailu, *Phys. Lett.* **B506** (2001) 207; W. D. Goldberger and M. B. Wise, [hep-th/0104170](#); R. Contino, L. Pilo, R. Rattazzi and E. Trincherini, [hep-ph/0108102](#).
- [35] N. Arkani-Hamed, T. Gregoire and J. Wacker, [hep-th/0101233](#).
- [36] G.v. Gersdorff and M. Quirós, work in progress.